hydrobiologia volume 323 (1996), contents

volume 323, no. 1, april 19, 1996

Use of an alternative method for monitoring total nitrogen concentrations in Florida lakes
by R.W. Bachmann & D.E. Canfield Jr. 1–8

Does the intermediate disturbance hypothesis comply with thermodynamics?
by S.E. Jørgensen & J. Padisák 9–21

Production and biomass of *Tenagomysis tasmaniae* Fenton, *Anisomysis mixta australis* (Zimmer) and *Paramesopodopsis rufa* Fenton from south-eastern Tasmania (Crustacea: Mysidacea)
by G.E. Fenton 23–30

Diet and predators of *Tenagomysis tasmaniae* Fenton, *Anisomysis mixta australis* (Zimmer) and *Paramesopodopsis rufa* Fenton from south-eastern Tasmania (Crustacea: Mysidacea)
by G.E. Fenton 31–44

Chydorus mutilus Kreis, 1921 – a postephippial form of *Chydorus sphaericus* (O. F. Müller, 1785)
by A. Brancelj 45–59

A comparison of solids collected in sediment traps and automated water samplers
by L.A. Bartsch, R.G. Rada & J.F. Sullivan 61–66

Influence of light on bacterioplankton in a tropical lake
by M. Lindell & H. Edling 67–73

Nature and occurrence of sulfoxidizing bacteria in Barégine developing in sulfurated thermal waters at Barèges (France)
by H. Hédoin, A. Couté, P. Kaiser & R. Laugier 75–81

volume 323, no. 2, april 26, 1996

Effects of macrophyte growth forms on invertebrate communities in saline lakes of the Wyoming High Plains
by W.M. Wollheim & J.R. Lovvorn 83–96

Determination of the trophic state by qualitative and quantitative phytoplankton analysis in two gravel pit lakes
by B. Tremel 97–105

The metabolism of organic matter in the hyporheic zone of a mountain stream, and its spatial distribution
by M. Pusch 107–118

Survival, osmoregulation and oxygen consumption of YOY coastal largemouth bass, *Micropterus salmoides* (Lacepede) exposed to saline media
by G.N. Susanto & M.S. Peterson 119–127

Decomposition rates and phosphorus concentrations of purple Loosestrife (*Lythrum salicaria*) and Cattail (*Typha* spp.) in fourteen Minnesota wetlands
by S.L. Emery & J.A. Perry 129–138

Description of *Phyllodiaptomus christineae* n.sp. from Thailand, and distinction of two subgenera within *Phyllodiaptomus* Kiefer, 1936 (Copepoda, Calanoida)
by H.J. Dumont, Y.R. Reddy & L.-o. Sanoamuang 139–148

Short note
How accurately can retention of benthic macrofauna by a particular mesh size be predicted from body size of organisms?
by T.A. Schlacher & T.H. Wooldridge 149–154

volume 323, no. 3, may 3, 1996

Effects of *Chironomus tentans* larval growth retardation on adult emergence and ovipositing success; implications for interpreting freshwater sediment bioassays
by K. Liber, D.J. Call, T.D. Dawson, F.W. Whiteman & T.M. Dillon 155–167
The biogeography of littoral *Lecane* Rotifera
by H. Segers 169–197

Invited papers section
Preface
by D.T. Crisp & R.H.K. Mann 199
Environmental requirements of common riverine European salmonid fish species in fresh water with particular reference to physical and chemical aspects
by D.T. Crisp 201–221
Environmental requirements of European non-salmonid fish in rivers
by R.H.K. Mann 223–235

Books received 237–238

Book review 239–242

Author index 243